

CHEST

THE CARDIOPULMONARY
AND CRITICAL CARE JOURNAL

FOR PULMONOLOGISTS, CARDIOLOGISTS, CARDIOTHORACIC SURGEONS,
CRITICAL CARE PHYSICIANS, AND RELATED SPECIALISTS

Inhaled Corticosteroids for Asthma Therapy: Patient Compliance, Devices, and Inhalation Technique

Mac G. Cochrane, MD; Mohan V. Bala, PhD; Kristen E. Downs, MSPH;
Josephine Mauskopf, PhD; and Rami H. Ben-Joseph, PhD
DOI: 10.1378/chest.117.2.542

This information is current as of November 15, 2005

The online version of this article, along with updated information and services, is located on the World Wide Web at:
<http://www.chestjournal.org/cgi/content/full/117/2/542>

CHEST is the official journal of the American College of Chest Physicians. It has been published monthly since 1935. Copyright 2005 by the American College of Chest Physicians, 3300 Dundee Road, Northbrook, IL 60062. All rights reserved. No part of this article or PDF may be reproduced or distributed without the prior written permission of the copyright holder. ISSN: 0012-3692.

AMERICAN COLLEGE OF
CHEST
P H Y S I C I A N S

Inhaled Corticosteroids for Asthma Therapy* Patient Compliance, Devices, and Inhalation Technique

Mac G. Cochrane, MD; Mohan V. Bala, PhD; Kristen E. Downs, MSPH;
Josephine Mauskopf, PhD; and Rami H. Ben-Joseph, PhD

Background: Patient compliance, inhalation devices, and inhalation techniques influence the effectiveness of inhaled medications.

Methods: This article presents the results of a systematic literature review of studies measuring compliance with inhaled corticosteroids, measuring inhalation technique with different inhalation devices, and estimating the proportion of inhaled drug that is deposited in the lung.

Results: Overall, patients took the recommended doses of inhaled medication on 20 to 75% of days. Frequency of efficient inhalation technique ranged from 46 to 59% of patients. Education programs have been shown to improve compliance and inhalation techniques. The lung deposition achieved with different inhalers depends on particle size as well as inhaler technique.

Conclusion: This review demonstrates that multiple factors may come between a prescription of an inhaled corticosteroid and the arrival of that medicine at its target organ, the lung.

(CHEST 2005; 117:542-550)

Key words: asthma; compliance; inhaler technique; inhalers

Abbreviations: DPI = dry powder inhaler; ICS = inhaled corticosteroid; MDI = metered-dose inhaler

Asthma is a common chronic inflammation of the airways that causes periodic attacks of wheezing and troubled breathing. Pharmacologic therapy is used to prevent and control asthma symptoms and reverse airflow obstruction. The major classes of asthma medications are β -agonists, methylxanthines, anticholinergics, leukotriene modifiers, inhaled corticosteroids, and corticosteroids (glucocorticoids). The majority of these medications are inhaled, and their effectiveness in clinical practice can be affected by many factors.

In this article, we examined three factors that appear to have an effect on the effectiveness of inhaled corticosteroid (ICS) treatment by conducting a review of the relevant literature. First, we

examined patient compliance with inhaled asthma therapy. Second, we examined inhalation technique. Third, we examined the impact of inhalation technique and inhalation device on drug deposition in the lungs. The main objective of the article was to summarize the research conducted so far into these three factors to provide information about the reliability of the inhaled route of administration of ICSs.

MATERIALS AND METHODS

The literature review started with a thorough key word search of the National Library of Medicine's MEDLINE database for articles on inhaled asthma therapy published between January 1986 and 1997. We examined the abstracts of these articles and identified articles that presented the results of studies of compliance, inhalation technique, and lung deposition of inhaled asthma drugs. We obtained these articles and examined their references for any other articles pertaining to these issues. We identified Australian and European articles from journals that are included in MEDLINE, such as *European Respiratory Journal*, *Medical Journal of Australia*, *Australian and New Zealand Journal of Medicine*, *European Journal of Respiratory Diseases*, and *Acta Allergologica*, as well as from international medical journals such as *Lancet*, *Asthma*, and *Respiratory Medicine*. From searching the references of the articles we obtained with the MEDLINE search, we found articles from European journals not listed in MEDLINE, such as *British Journal of Disease of the Chest*. In

CHEST / 117 / 2 / FEBRUARY, 2000 543

Table 2—Proper Inhalation Technique*

Study	No.	Year	Trained	Type of Inhaler	Physician Assessment, %		Frequency of Patient Technique, %
					"Good"	"Inadequate"	
Dowling et al. ¹	41	1992	Yes	Rotahaler	5	95	27
Hilton ²⁰	62	1990	NS	Rotahaler	50	50	21
Pedersen et al. ²¹	39	1996	Yes	Rotahaler	45	55	30
Hilton ²⁰	262	1990	NS	MDI	45	55	30
Crompton ²²	132	1996	Yes	MDI	6	94	54
Pedersen et al. ²¹	86	1995	No	MDI	46	54	15
Honley and Bailey ²³	86	1995	Yes	MDI	86	14	8
Honley and Bailey ²³	321	1976	Yes	MDI	28	72	59
Pedersen and Crompton ²⁴	69	1990	No	MDI	52	48	37
Cayard and Oreluck ²⁵	46	1990	Yes	MDI	58	42	22
Wilton ²⁶	35	1990	NS	Spacer device	78	22	4
Pedersen et al. ²¹	85	1996	Yes	Rotahaler	54	46	19
Hilton ²⁰	23	1990	NS	Rotahaler	54	46	19

*NS = not specified.

†The study's categories were reclassified as follows: failless = good; doubtful or adequate = adequate; totally inadequate or probably inadequate = inadequate.

‡The study's categories were reclassified as follows: efficient = good; doubtful = adequate; inefficient = inadequate.

§The study's categories were reclassified as follows: correct = good; doubtful = adequate; incorrect = inadequate.

Melchor et al.²⁵ showed an increase in peripheral lung deposition of salbutamol, but not in total lung deposition, with the use of a spacer device.

Proper inhalation technique can also significantly affect the amount of drug delivered to the lung. Newman et al.²⁶ showed that for patients who could not coordinate inhalation and actuation at baseline (44% of the population), instruction improved lung deposition from 7.2 to 22.8%. Furthermore, before instruction, the patients who coordinated inhalation and actuation had a higher lung deposition (18.6%) than those who did not (7.2%).

Discussion

In this paper, we examined three factors influencing the effectiveness of asthma treatment with ICSs in practice—patient compliance, inhalation technique, and lung aerosol deposition. We found that only a small percentage of the prescribed dose of an ICS is likely to reach the target organ, the lung, because of patient noncompliance with the prescribed dose, difficulty in correct use of the inhaler, and the ability of a properly used inhaler to deliver the drug to the lung.

As with any chronic disease, patient compliance is an important determinant of therapeutic success. Greer and Levstek²⁷ divide the factors that are correlated with noncompliance into four categories: patient variables, interactions between

physician or medical staff and patients or family, medication characteristics, and nature of asthma. Our evaluation indicates that patients were more likely to underuse ICSs than overuse them. Some examples of factors related to noncompliance and inhaled drugs include patients taking long-term medication who stop treatment if they have not experienced an attack for an extended period.²⁸ The delayed clinical impact of ICSs compared with bronchodilator drugs may be an additional factor in noncompliance. The possibility of psychosocial issues in noncompliant patients should not be overlooked. Bosley et al.²⁹ found in a prospective cohort that noncompliant patients had a significantly higher prevalence of depression than compliant patients. Noncompliant patients also had a higher, although not statistically significant, prevalence of anxiety.

There is conflicting evidence about the relationship between the number of inhalations prescribed per day and compliance rates. Two published studies supported the assumption that there is an inverse relationship between the number of inhalations prescribed per day and compliance.^{17,41} However, this inverse relationship was not supported in studies by Bosley et al.²⁹ and Tonggood et al.⁴²

Education has been shown to increase compliance with dosing regimens. The study by van der Walen et al.¹⁴ showed this using electronic monitoring for compliance. A recent meta-analysis of the effects of

Table 3—Observed Errors in Inhalation Checklists*

Factors	MDI									
	Pedersen et al. ²¹	Pedersen et al. ²¹	van Berendse et al. ²²	van Berendse et al. ²²	Manzella Epstein et al. ²³	Honley et al. ²³	Smith et al. ²⁴	Scott et al. ²⁵	Nimmo et al. ²⁶	Cayard and Oreluck ²⁵
Number of patients	132	851	56	57	34	68	84	59	19	69
Remove cap, %	49	34	57	57	32	18	32	18	10	32
Shake inhaler, %			2			12		0	0	
Hold inhaler upright, %						66				
Place mouthpiece of inhaler into one end of spacer, %						19				
Position inhaler with index finger on top and thumb supporting the bottom, %										
Exhale before filling, %	45	51	66			58	0	15	26	75
Tilt head back, %	12	14				15	30			
Place spacer tube between teeth, %						25	13		0	
Close mouth, %						16				
Release medication, %	55	17	68			34	40	25	42	47
Begin to breathe in and then activate inhaler, %	67	29	70			43	66	45	40	47
Inhale slowly and deeply, %	42	39	53			31	75	59	25	32
Hold breath, %						49		20	32	
Breathe out through nose, %						54	23	46	35	NA
Breathe released slowly, %										
Wait before taking next puff, %										

*Other Inhalers

Factors	Other Inhalers									
	Pedersen et al. ²¹	van Berendse et al. ²²	van Berendse et al. ²²	Dowling et al. ¹	Nimmo et al. ²⁶	1st Puff	2nd Puff	1st Puff	2nd Puff	Wilton ²⁶
Number	39	257	257	41	14	14	14	14	14	144
Type of Inhaler	Rotahaler	Mix of dry powder inhalers		Rotahaler	Rotahaler	Rotahaler	Rotahaler	Rotahaler	Rotahaler	Turbuhaler
Cover removed, %	15	14	14	0	0	0	0	0	0	2
Loading drug substance, %				51	15	15	15	15	15	3
Hold inhaler at correct angle, %	46	7	7	0	0	0	0	0	0	1
Rotation sequence correct, %	19									3
Air fully inhaled, %	62	66	66	83	0	0	0	0	0	3
Exhale before filling, %	13			17	15	15	15	15	15	0
Tilt head back, %	36	4	4							0
Drug capsule not emptied, %										0
Place mouthpiece between lips, %										0
Release medication, %	41	10	10	29	23	23	23	23	23	0
Inhale forcefully and deeply, %	49	53	53	76	83	83	83	83	83	4
Hold breath, %										0
Breathe out through nose, %										0
Breathe released slowly, %										0
Wait before repeating step, %										0

*If blank, error was not reported. NA = not applicable for second puff.

†After instruction, others are preinstruction.

psychoeducational care in adults with asthma by DeVine²⁷ also showed improvements in self-reported and physician-assessed compliance associated with education programs.

Other routes of administration for anti-asthma drugs may improve compliance. Kelloway et al.¹⁴ found that compliance was higher with prescribed oral medications compared with inhaled anti-inflam-

matory medications for asthma. However, their study was a retrospective database study that compared oral theophylline with two inhaled anti-inflammatory medicines. Thus, in their study, higher compliance with the oral drug might have been related to more rapid impact of theophylline on symptoms rather than the mode of administration. It is also important to note that oral medications are not affected by two of the three factors discussed in this paper that may reduce drug effectiveness: inhalation technique and lung deposition. Well-designed prospective studies comparing oral and inhaled asthma therapies could help further to determine the importance of route of administration.

Evidence suggests that noncompliance has an effect on patient outcomes. For example, Horn et al⁴⁸ treated 160 patients for as long as 9 months with increasing doses of inhaled salbutamol and beclomethasone dipropionate. The authors found that patients whose conditions improved had higher urinary salbutamol concentrations compared with those with persistent problems.

The effectiveness of inhaler therapy also depends on the inhaler technique. Patients may not be adequately instructed in inhaler technique, thereby reducing the amount of drug delivered to the lungs.³⁰ Cochrane⁴⁹ states that "it is important to reinforce the simple concept that failure to instruct patients on how to use inhalers and to reinforce these instructions will decrease compliance, whatever the drug or inhalation device." Our study found that subjective assessments of inha-

tion technique by physicians provided a wide range of values in patients using the inhaler properly. Technique is highly dependent on the type of inhaler. In our evaluation, we found that patients using a Rotahaler and an MDI had similar rates of "inadequate" technique. However, patients using DPIs other than the Rotahaler, such as the Turbuhaler and the Diskhaler, had lower rates of "inadequate" technique. DPIs... have the intrinsic advantage that there is a natural coordination between generation of the aerosol cloud and inspiration.⁴⁶ The lack of need for coordination of two activities by the patient using a DPI makes it more likely that their inhalation technique with these devices will be good. As expected, patient groups using a spacer in addition to an MDI had lower rates of "inadequate" technique than the majority of patient groups using an MDI alone.

Because physician rating of patient technique is subjective, other methods of evaluating patient technique are needed, particularly so that the measurement scale can be consistent across studies. The study by Goodman et al⁵⁰ is an example of a study using an objective technique for measuring inhaler skills, a computer sensing device.

Several studies have shown that education can have a large impact on the percentage of patients who use an inhaler correctly.^{51,52} The study by Goodman et al⁵⁰ using a computer sensing device, indicated that women may have a greater need for education on inhaler technique because of less ef-

Table 4—Drug Delivered to Target Area: MDI*

MDI		MDI With Spacer	
Study	Drug MMD, μ m	Study	Drug MMD, μ m
Dolovich et al ¹² (n = 127)	Label 2.21	Newman et al ¹³ (n = 9)	Label 3.2
Newman et al ¹³ (n = 9)	Label 3.21	Melcher et al ¹⁴ (n = 91)	Salbutamol 2.5-3.9
Newman et al ¹⁴ (n = 19)	Salbutamol		
Melcher et al ¹⁵ (n = 217)	Salbutamol 2.5-2.9		
Zainudin et al ¹⁶ (n = 9)	Salbutamol + label 2.3		
Bongers et al ¹⁷ (n = 13)	Turbuhaler, not specified		
Range		Range	
	14		21
	9		19-21
	7-23		
	10-13		
	19-22		
	11		
	8		
	7-23		19-21

*MMD = mass median diameter for inhaled particle.

†Based on a maximum of 6.8% delivered to the right lung.

‡Label refers to a radioisotope attached to a nontherapeutic particle.

§For patients who could not coordinate inhalation and actuation, before and after instruction.

¶For patients who could coordinate inhalation and actuation, before and after instruction.

‡Range for asthma patients in healthy subjects.

Table 5—Drug Delivered to Target Area: Inhalers*

Breath-Actuated Inhaler		DPI	
Study	Drug MMD	Study	Drug MMD, μ m
Newman et al ¹⁴ (n = 19)	Salbutamol, not specified	Melcher et al ¹⁵ (n = 29)	Salbutamol 2.7-3.0
		Zainudin et al ¹⁶ (n = 9)	Salbutamol + label 2.3
		Bongers et al ¹⁷ (n = 10)	Budonide 2.3-3.0
		Bongers et al ¹⁷ (n = 13)	Turbuhaler, not specified
Range		Range	
			19-22
			9-28

*MMD = mass median diameter for inhaled particles.

†Range for patients who could coordinate inhalation and actuation and those who could not after all were trained to use the inhaler.

‡Range for asthma patients and healthy subjects.

§Label refers to a radioisotope attached to a nontherapeutic particle.

¶Range for slow to fast inspiratory flow.

‡Range based on inhaled drug dose.

fective use of the inhaler. They suggest that this problem with inhaler technique might explain why women experience more severe asthma symptoms than men. However, a recent study in a health maintenance organization population by Osborne et al⁴⁷ showed that women had more severe asthma symptoms but that their MDI technique was no different from that of the men when skills were ranked using a 10-item checklist. Osborne et al⁴⁷ suggest one possible reason for the difference in results could be that the electronic sensor is a more accurate measure of inhaler skills than the 10-item checklist.

Studies have shown that written instructions are not sufficient and that verbal instructions and demonstrations and practice sessions need to be included.⁵⁰ Simple teaching devices can be built to develop coordination skills needed with MDIs.⁴⁸ Patients also need to be checked periodically to make sure that their skills have not eroded over time.⁵⁰

The goal of inhaled therapy is to deliver medication directly to the lungs. The patient is at an advantage if the amount of the drug reaching the lungs is maximized and the amount deposited on the oropharyngeal region is minimized. The likelihood of adverse events can be reduced by reducing the amount of drug that reaches the oropharynx.⁴⁹ Many factors affect the amount of drug that reaches the lung, including inhaler technique and inhaler type, fine particle dose, and particle distribution. Of the studies in our analysis, only about 7 to 23% of any drug delivered by an MDI enters the respiratory tract. The rest of the drug is either deposited in the oropharynx or swallowed, leading to potential systemic side effects. Studies have shown that the amount of drug delivered to the

lungs can be almost doubled and that the amount of drug deposited in the oropharynx greatly reduced through the use of large-volume spacers.¹⁵ A study using the Turbuhaler showed an increased in drug deposition in the lung.³⁷ Jackson and Lipworth¹³ report two studies that have shown that the use of a Turbuhaler is associated with greater efficacy of ICSs. Thus, the increased deposition might be associated with increased drug efficacy.

The lung deposition studies reported in our article show mixed results and do not clearly demonstrate that one type of inhaler is superior to another for use with ICSs. There is a need for more research in this area. Lung deposition depends critically on particle dynamics and jet flow,⁵⁰ which vary with drug formulation, inhaler type, and patient inhaler technique. Large crossover studies will need to be designed that control for drug type and particle size as well as inhaler technique. The outcomes of such studies should be both direct measures of drug deposition as well as measures of patient outcomes.

A key to efficacy for ICSs for treatment of asthma is the amount of drug prescribed that reaches the target organ. This review of patient compliance, inhalation technique, and inhaler type demonstrates the importance of these three factors in the appropriate delivery of ICSs to the lung. Health-care providers should consider these factors when evaluating and prescribing asthma medications.

REFERENCES

- Haynes RR, Taylor DW, Sackett DL, eds. Compliance in health care. Baltimore, MD: Johns Hopkins University Press, 1979

- 2 Cochrane GM. Compliance and outcomes in patients with asthma. *Drugs* 1996; 52:12-19
- 3 Bosley CM, Parry DT, Cochrane GM. Patient compliance with inhaled medication: does combining beta-agonists with corticosteroids improve compliance? *Eur Respir J* 1994; 7:504-509
- 4 Rand CS, Wise RA. Measuring adherence to asthma medication regimens. *Am J Respir Crit Care Med* 1994; 149:569-576
- 5 Braunstein GL, Trinquet G, Harper AE, et al. Compliance with nedocromil sodium and a nedocromil sodium/salbutamol combination. *Eur Respir J* 1996; 9:882-898
- 6 Murphy S. Discussion section. *Am J Respir Crit Care Med* 1994; 149:577-578
- 7 Dompeling E, Van Grunven PM, Van Schayck CP, et al. Treatment with inhaled steroids in asthma and chronic bronchitis: long-term compliance and inhaler technique. *Fam Pract* 1992; 9:161-166
- 8 Young M, O'Connor SA, Parry DT, et al. Compliance with prescribed drug therapy in asthma. *Respir Med* 1994; 88:31-35
- 9 Rand CS, Wise RA, Nides M, et al. Metered-dose inhaler adherence in a clinical trial. *Am Rev Respir Dis* 1992; 146:1599-1594
- 10 Coutts JAP, Gibson NA, Paton JY. Measuring compliance with inhaled medication in asthma. *Arch Dis Child* 1991; 67:332-333
- 11 Manzella B, Brooks C, Richards J, et al. Assessing the use of metered dose inhalers by adults with asthma. *J Asthma* 1993; 26:223-230
- 12 Dolovich M, Ruffin RE, Roberts R, et al. Optimal delivery of aerosols from metered dose inhalers. *Chest* 1981; 80(suppl): 911-915
- 13 Jackson C, Lipworth B. Optimizing inhaled drug delivery in patients with asthma. *Br J Gen Pract* 1995; 45:683-687
- 14 Doolley CT, Paterson JW, Connolly ME. Clinical pharmacology of beta-receptor-blocking drugs. *Clin Pharmacol Ther* 1989; 10:765-799
- 15 Chervinsky H. Is total particle dose more important than particle distribution? *Respir Med* 1997; 91(suppl A):17-19
- 16 van der Palen J, Klein J, Rivers M. Compliance with inhaled medication and self-treatment guidelines following a self-management programme in adult asthmatics. *Eur Respir J* 1997; 10:652-657
- 17 Mann MC, Eliasson O, Patel K, et al. An evaluation of severity-modulated compliance with p.i.d. dosing of inhaled beclomethasone. *Chest* 1992; 102:1342-1346
- 18 Mawhinney H, Spector SL, Kirsman RA, et al. Compliance in clinical trials of two nonbronchodilator, antiasthma medications. *Ann Allergy* 1991; 66:294-299
- 19 Ajfer A, Reisman S, Liffick C, et al. Adherence with twice-daily dosing of inhaled steroids: socioeconomic and health-related differences. *Am J Respir Crit Care Med* 1998; 157:1310-1317
- 20 Hilton S. An audit of inhaler technique among asthma patients of 34 general practitioners. *Br J Gen Pract* 1990; 40:505-506
- 21 Pedersen S, Frost L, Arnfred T. Error in inhalation technique and efficiency in inhaler use in asthmatic children. *Allergy* 1996; 41:118-124
- 22 Crompton GK. Nebulized or intravenous beta₂-adrenoceptor agonist therapy in acute asthma? *Eur Respir J* 1990; 3:125-126
- 23 Horsley MC, Balle CR. Risk factors for inadequate use of pressurized aerosol inhalers. *J Clin Pharm Ther* 1988; 13:139-143
- 24 Paterson M. Crompton C. Use of pressurized aerosol by asthmatic patients. *BMJ* 1976; 272:76-77
- 25 Gayard P, Orsick J. Manual utilization of aerosol devices for asthma. *Respiration* 1990; 40:47-52
- 26 van Reccondonk I, Mesters I, Mulder A, et al. Assessment of the inhalation technique in outpatients with asthma or chronic obstructive pulmonary disease: using a metered-dose inhaler or dry powder device. *J Asthma* 1995; 32:273-279
- 27 Epstein SW, Manning CPR, Ashley MJ, et al. Survey of the clinical use of pressurized aerosol inhalers. *Can Med Assoc J* 1976; 120:513-516
- 28 Scott-Smith W. Sterning inhaler technique in general practice. *Practitioner* 1986; 230:96
- 29 Nimmo CJ, Renner D, Chen NM, et al. Assessment of patient acceptance and inhalation technique of a pressurized aerosol inhaler and two breath-actuated devices. *Ann Pharmacother* 1993; 27:922-927
- 30 Goodman D, Israel E, Rosenborg M, et al. The influence of age, diagnosis, and gender on proper use of metered-dose inhalers. *Am J Respir Crit Care Med* 1994; 150:1256-1261
- 31 Watson JBC. The acceptability and efficacy of terbutaline given by metered-dose, powder inhalation ("Breathe Turbulator") in asthmatic hospital out-patients: a multi-center study. *Curr Med Res Opin* 1990; 11:651-660
- 32 Bailey W, Richard J, Brooks M, et al. A randomized trial to improve self-management practices of adults with asthma. *Arch Intern Med* 1990; 150:1664-1668
- 33 Newman SP, Miller AB, Leonard-Jones TL, et al. Improvement of pressurized aerosol deposition with Nebulizer spacer device. *Thorax* 1994; 39:935-941
- 34 Newman SP, Weisz AWB, Talanc N, et al. Improvement of drug delivery with a breath-actuated pressurized aerosol for patients with poor inhaler technique. *Thorax* 1991; 46:712-716
- 35 Mclodur R, Riddiscombe MF, Mak VHF, et al. Lung deposition patterns of directly labeled salbutamol in normal subjects and in patients with reversible airflow obstruction. *Thorax* 1983; 46:506-511
- 36 Zainudin RMZ, Riddiscombe M, Tolfree SF, et al. Comparison of bronchodilator responses and deposition patterns of salbutamol inhaled from a pressurized metered dose inhaler, as a dry powder, and as a nebulized solution. *Thorax* 1980; 45:469-473
- 37 Borgström L, Derrin E, Ståhl E, et al. The inhalation device influences lung deposition and bronchodilating effect of terbutaline. *Am J Respir Crit Care Med* 1996; 153:1636-1640
- 38 Borgström L, Bondesson E, Morén F, et al. Lung deposition of budesonide inhaled via Turbuhaler: a comparison with technical sulphate in normal subjects. *Eur Respir J* 1994; 7:89-93
- 39 Greer TL, Lesick D. Medication compliance and asthma: overlooking the tree because of the forest. *J Asthma* 1996; 33:203-211
- 40 Bosley CM, Frobey JA, Cochrane GM. The psychological factors associated with poor compliance with treatment in asthma. *Eur Respir J* 1995; 8:899-904
- 41 Malo JL, Cartier A, Ghezzo H, et al. Comparison of four-times-a-day and twice-a-day dosing regimens in subjects requiring 1200 µg or less of budesonide to control mild to moderate asthma. *Respir Med* 1995; 89:537-543
- 42 Toogood J, Baskerville C, Jennings B, et al. Influence of dosing frequency and schedule on the response of chronic asthmatics to the aerosol steroid, budesonide. *J Allergy Clin Immunol* 1992; 70:285-298
- 43 Devine E. Meta-analysis of the effects of psychosocial factors on adherence to asthma therapy. *Respir Med* 1996; 140:1-10
- 44 Kellway JS, Wyatt RA, Adis SA. Comparison of patients' compliance with prescribed oral and inhaled asthma medications. *Arch Intern Med* 1994; 154:1349-1352
- 45 Horn GR, Clark TJH, Cochrane GM. Compliance with inhaled therapy and morbidity from asthma. *Respir Med* 1990; 84:67-70
- 46 Gauderton D. General factors influencing drug delivery to the lung. *Respir Med* 1997; 91(suppl A):13-16
- 47 Osborne M, Vollmer W, Linton K, et al. Characteristics of patients with asthma within a large HMO: a comparison by age and gender. *Am J Respir Crit Care Med* 1998; 157:123-125
- 48 Lee H, Evans H. Aerosol inhalation teaching device. *J Pediatr* 1987; 110:249-252
- 49 Barnes PJ, Pedersen S. Efficacy and safety of inhaled corticosteroids in asthma. *Am Rev Respir Dis* 1992; 145:511-526
- 50 Kim C, Trujillo D, Sackner M. Size aspects of metered-dose inhaler aerosols. *Am Rev Respir Dis* 1985; 132:137-142

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ **BLACK BORDERS**
- ☐ **IMAGE CUT OFF AT TOP, BOTTOM OR SIDES**
- ☐ **FADED TEXT OR DRAWING**
- ☐ **BLURRED OR ILLEGIBLE TEXT OR DRAWING**
- ☐ **SKEWED/SLANTED IMAGES**
- ☐ **COLOR OR BLACK AND WHITE PHOTOGRAPHS**
- ☐ **GRAY SCALE DOCUMENTS**
- ☐ **LINES OR MARKS ON ORIGINAL DOCUMENT**
- ☐ **REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY**
- ☐ **OTHER:** _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.